

# Smarter Balanced Assessment Consortium:

## **Practice Test Scoring Guide**

Grade 11


Published August 19, 2013

Prepared by the American Institutes for Research®




© Smarter Balanced Assessment Consortium, 2013 Descriptions of the operation of the Test Delivery System, Test Information Distribution Engine, and related systems are property of the American Institutes for Research® (AIR) and are used with permission of AIR.





• correct placement of the graph with its vertex at (4, 2)






Consider the function  $f(x) = x^2 - 5x - 14$ . Which of the numbers in the chart are zeros of the function? Select Yes or No in each row.

Is this a zero of the function?	Yes	No
2		
7		
-2		
-7		


For this item, a full-credit response (2 points) includes:

- a check in the "No" column for 2 AND
- a check in the "Yes" column for 7 AND
- a check in the "Yes" column for -2 AND
- a check in the "No" column for -7

For partial credit (1 point), the student correctly checks at least 3 boxes.




• the value  $\frac{1}{2}$  in the box



Create the function for the graph.

For this item, a full-credit response (1 point) includes:

• a function equivalent to f(x) = (x - 2)(x - 2)(x + 1)(x + 3)(x - 4) by the commutative property



• the symbol  $\leq$  inside the box

#### 676

A car rental company charges customers an initial charge plus a daily charge to rent cars. The initial charge is \$30 and the daily charge is \$25.

The rental company charged Jacob \$180.

Create an equation that can be used to find the number of days, x, Jacob rented the car.

Click the buttons to create your answer.


$\bullet \bullet \bullet \bullet \bigotimes$
123 x
$456 + - \times \div$
789 < ≤ = ≥ >
0 Η Π°Ο ΙΙ π i
sin cos tan arcsin arccos arctan

For this item, a full-credit response (1 point) includes:

• a correct equation, such as 25x + 30 = 180

The table shows several inputs and outputs for two functions, fand g, that are both continuous on the interval 0 to 8.

x	<i>f</i> ( <i>x</i> )	g(x)
0	-5	120
1	-4	103
2	-1	86
3	4	69
4	11	52
5	20	35
6	31	18
7	44	1
8	59	16



There is exactly one solution for which f(x) = g(x).

Click the number line to show the unit interval for x in which the solution to f(x) = g(x) must lie.

For this item, a full-credit response (1 point) includes:

• the interval [5, 6] selected

The \$1000 prize for a lottery is to be divided evenly among the winners. Initially there are x winners, but then one more winner comes forward, causing each winner to receive \$50 less.


### **681**

Create an equation that represents the situation and can be used to solve for x, the initial number of winners.

$\bullet \bullet \bullet \bullet \bullet \\ \blacksquare$
123 x
$4 5 6 + - \times \div$
$789 < \leq = \geq >$
0 Η [] ()    √] <sup>[</sup> √] π i
sin cos tan arcsin arccos arctan

For this item, a full-credit response (1 point) includes:

• a correct equation, such as  $\frac{1000}{x} = \frac{1000}{(x+1)} + 50$ 




$$\sqrt{64} = x - 3 \qquad \sqrt{39} - 3 = x$$
$$x - \sqrt{5} = \sqrt{20} \qquad \sqrt{3x} = 75$$
$$\sqrt{x} = \frac{\sqrt{16}}{8} \qquad 2x = \sqrt{100}$$

• 
$$\sqrt{64} = x - 3$$
  
AND

• 
$$\sqrt{3x} = 75$$
  
AND

• 
$$2x = \sqrt{100}$$



- a and c having opposite signs and b being any value AND
- a and c having the same sign and b being any value OR when a is zero and c is not zero and b being any value AND
- *c* being zero and *a* not being zero and *b* being any value

For example,

- y = 3|x + 2| + (-5)AND
- y = 3|x + 2| + 5AND
- y = 3|x + 2| + 0


For partial credit, the student completes each task for 1 point each.



The graphs of y = f(x) and y = g(x) are shown.

Drag points onto the coordinate grid to show


- a solution for y = f(x),
- a solution for y = g(x), and
- a solution for f(x) = g(x).




For this item, a full-credit response (2 points) includes:

- "Solution f(x)" on the line of f(x)
  AND
- "Solution g(x)" on the line of g(x)
  AND
- "Solution f(x) = g(x)" on the intersection of line f(x) and line g(x)

For partial credit (1 point), the student places "Solution f(x) = g(x)" correctly or the student places both of the other points correctly.



• three points correctly plotted on the curve of  $y = \frac{3}{\sqrt{x}}$ 



For example,

682	

Consider triangle ABC, where angle C is a right angle.


Drag possible measures of angle A into the correct column.

$\cos A < \sin A$	$\cos A = \sin A$	$\cos A > \sin A$
Poss	ible Measures of An	_
5° 25°	35°	15° 45°
55°	65° 75°	45 85°

For this item, a full-credit response (2 points) includes:

- 55°, 65°, 75°, and 85° in the "cosA < sinA" column AND
- $45^{o}$  in the "cosA = sinA" column AND
- $5^{o}$ ,  $15^{o}$ ,  $25^{o}$ , and  $35^{o}$  in the "cosA > sinA" column

For partial credit (1 point), the student correctly fills out 2 columns.



- a = 3 and  $b \neq 1$ AND
- a = 3 and b = 1AND
- *b* = 5


For partial credit, each correct task is worth 1 point.

202	1 THL	<b>Delete</b>	×		
A teacher has a classroom with 30 boys and girls. The class has both 9th and 10th graders.			Stude	ents by Gender	r and Grade
The probability of a randomly			Gender	Grade 9	Grade 10
selected student from the class			Female		
being a female <b>or</b> in grade 9 is		Female			
equal to $\frac{19}{30}$ .					
Drag tick marks into each		Male			
section of the chart to show					
how this can be true.					

- 11 tally marks in the "Grade 10 Male" box AND
- a total of 19 tally marks in the "Grade 10 Female," "Grade 9 Male," and "Grade 9 Female" boxes combined


For example,

Gender	Grade 9	Grade 10
Female	ШТ I	JHT 11
Male	ШТ I	ШТ ШТ I



- a point at (2,3) AND
- a point at (6, 5)


For partial credit (1 point), the student correctly plots the translation of point *A* or point *B*.



- the gold star at the point (7,5) AND
- the red diamond at the point (8.75, 0)

For partial credit (1 point), the student correctly places the star or the diamond.

74



For this item, a full-credit response (1 point) includes:

• a vertical line at 63 inches



Melissa and Carrie both drew right triangles. The length of the hypotenuse in each triangle is  $\sqrt{130}$  units.

The perimeter of Melissa's triangle is  $14 + \sqrt{130}$  units.

A. Use the Connect Line tool to draw Melissa's triangle.

The perimeter of Carrie's triangle is  $16 + \sqrt{130}$  units.

B. Use the Connect Line tool to draw Carrie's triangle.

🕟 🕞 Delete 🛛 🗙 Add Point 🛶 🏶 Connect Line 🛁		
Melissa's Triangle	Carrie's Triangle	
1 unit	1 unit	

For this item, a full-credit response (2 points) includes:

- a right triangle with legs of 3 units and 11 units under "Melissa's Triangle" AND
- a right triangle with legs of 7 units and 9 units under "Carrie's Triangle"


For partial credit, the student completes either task for 1 point.



A ball is thrown in the air. The height of the ball in terms of time is modeled by the graph shown.

A second ball is thrown from a lower initial height and reaches a higher maximum height.


- Select an equation that represents the height of the second ball in terms of time.
- Use the Add Point tool to plot two points on the grid: the initial height of the second ball and its maximum height.



For this item, a full-credit response (2 points) includes:

- the equation  $y = -x^2 + 5x + 3$ AND
- a point at (0,3) and (2.5,9.25)

For partial credit, the student completes the above tasks for 1 point.



• the expression  $x^3 + 3x^2 + \frac{\pi x^3}{12}$ 

#### 685

The functions  $f(x) = 500(1.015)^x$  and  $g(x) = 500(1.021)^x$  give the total amounts in two different savings accounts after x years.

How do the graphs of f(x) and g(x) compare?


- A They have the same y-intercept, but the graph of f(x) rises more quickly over time.
- **(B)** They have the same *y*-intercept, but the graph of g(x) rises more quickly over time.
- © The function f(x) has a greater y-intercept and rises more quickly over time.
- **(b)** The function g(x) has a greater *y*-intercept and rises more quickly over time.

For this item, a full-credit response (1 point) includes:

• option B



The graph of exponential function f(x) is shown.



What is the value of f(6)?



For this item, a full-credit response (1 point) includes:

• the value 64